arXiv:1202.1139v1 [math.CO] 6 Feb 2012

André permutations of the second kind associated to strictly binary increasing trees and left to right minima in their sub-permutations

Filippo Disanto[*]

February 7, 2012

Abstract

The present work concerns André permutations of the second kind which are associated to strictly binary increasing tree. In particular we focus on their sub-permutations finding a bivariate generating function which provide their enumeration with respect to the size and to the number of left to right minima.

1 Introduction

The aim of this paper is to link three main concepts. The first is the one of *André* permutations of the second kind which have been introduced in [5] as permutations related to *binary increasing* trees. The second is the one of *strictly* binary increasing trees which have been considered for example in [3] and the third is the one of *sub-permutations*. This last tool has been used to define class of permutations such as *simsun* permutations, see [4] and the related references for enumerative properties concerning permutations belonging to this class.

What we want to consider in this work is indeed the class of sub-permutations of those André permutations of the second kind which are related to strictly binary increasing trees.

In Section 3.1 we will provide their enumerations with rispect to the size and in Section 3.2 we will count them with respect to the size and to the number of left to right minima. Moreover we will see how, concerning the class of permutations we will consider, the set of left to right minima corresponds to a particular path of the corresponding binary increasing trees.

---

[*] Institut für Genetik, Universität zu Köln.



Figure 1: The five trees in $\mathcal{B}_4$.

2 Definitions

The set of permutations of size n will be denoted by $\mathcal{S}_n$. If $\pi = (\pi_1\pi_2 ... \pi_n) \in \mathcal{S}_n$ the set of its *left to right minima* is denoted by $L(\pi)$ and its elements are those entries π_i such that if $j < i$, then $\pi_i < \pi_j$.

A *binary increasing* tree is a rooted (non planar) tree such that its nodes have outdegree $0, 1$ or 2; nodes with outdegree 0 are also called the *leaves* of the tree. Moreover, for such a tree we require that the set of nodes is totally ordered by labels in such a way going from the root to any leaf we always find an increasing sequence of numbers. In particular, if x and y are two nodes of such a tree, we will write $x \prec y$ when the label of x is less than the label of y. The set of binary increasing trees will be denoted by $\mathcal{B}$ while we will use the symbol $\mathcal{B}_n$ to denote the subset of $\mathcal{B}$ made of those trees with n nodes.

According to [5] a tree in $\mathcal{B}$ can be drawn in the plane in a *unique* way respecting the following two conditions $a1$) and $a2$):

$a1$) if a node has only one child then this child is drawn on the right of its direct ancestor,

$a2$) if a node x has two children y and z with $y \prec z$, then y is drawn on the left of x while z on the right.

In Fig. 1 we have depicted, respecting the previous two conditions, the trees in $\mathcal{B}_4$.

In [5] the authors define the the set of *André* permutations of the second kind $\mathcal{A}_n$ which is a subset of $\mathcal{S}_n$ equinumerous to $\mathcal{B}_n$. Indeed consider the following procedure $\phi : \mathcal{B}_n \to \mathcal{S}_n$:

1) given $t \in \mathcal{B}_n$ draw t according to $a1$) and $a2$);
2) each leaf collapse into its direct ancestor whose label is then modified receiving on the left the label of the left child (if any) and on the right the label of its right child. We obtain in this way a new tree whose nodes are labelled with sequences of numbers;
3) starting from the obtained tree go to step 2).

The algorithm ϕ ends when the tree t is reduced to a single node whose label is then a permutation $\phi(t)$ of size n.

The set $\mathcal{A}_n$ is defined as $\{\phi(t) \in \mathcal{S}_n : t \in \mathcal{B}_n\}$. Looking at Fig. 1 the corresponding permutations in $\mathcal{A}_4$ are (from left to right) $(3214), (1234), (2134), (1324)$ and (2413).

André permutations, as binary increasing trees, are enumerated with respect to the size by the so called *Euler* numbers whose exponential generating function is:

$$A(x) = \sec(x) + \tan(x) - 1.$$

The first terms of the sequence are: $1, 1, 2, 5, 16, 61, 272, 1385, 7936, ...$ and they corresponds to entry $A000111$ in [7].

A tree in $\mathcal{B}$ which does not have nodes of outdegree 1 is said to be *strictly* binary. We will denote the set of these trees having size n by the symbol $\tilde{\mathcal{B}}_n$. Similarly, the subset of $\mathcal{A}_n$ made of those permutations corresponding to strictly binary increasing trees will be denoted by $\tilde{\mathcal{A}}_n$ and we also define $\tilde{\mathcal{A}} = \bigcup_i \tilde{\mathcal{A}}_i$.

Given a permutation $\pi \in \mathcal{S}_n$ a *sub-permutation* of π is a permutation obtained from π by considering only the entries $1, 2, ..., k$ with $1 \leq k \leq n$. If σ is a sub-permutation of π we will write $\sigma \lhd \pi$. Sub-permutations are used in the literature to define classes of permutations. As an example we can consider the so called *simsun* permutations, studied for example in [4]. A permutation is said to be *simsun* if each sub-permutation has no double descent.

Now we are ready to define the object of our work. In what follows we will study the sets

$$sub_n(\tilde{\mathcal{A}}) = \{\sigma \in \mathcal{S}_n : \exists\, \pi \in \tilde{\mathcal{A}}\ \sigma \lhd \pi\},$$

$$sub(\tilde{\mathcal{A}}) = \bigcup_i sub_i(\tilde{\mathcal{A}}).$$

We want to observe that in general $sub_n(\tilde{\mathcal{A}}) \nsubseteq \mathcal{A}_n$. Indeed we have that $3241 \lhd 32415 \in \tilde{\mathcal{A}}$ but $3241 \notin \mathcal{A}_4$.

Finally we define $sub_{n,m}(\tilde{\mathcal{A}})$ which is composed by those permutations π in $sub_n(\tilde{\mathcal{A}})$ having $|L(\pi)| = m$; the cardinality of $sub_{n,m}(\tilde{\mathcal{A}})$ will be found in Section 3.2.

3 Enumerations

In this section we provide the enumerations of $sub_n(\tilde{\mathcal{A}})$ and $sub_{n,m}(\tilde{\mathcal{A}})$ using the existing correspondance between permutations and binary increasing trees. In particular, in Section 3.2 we will reach our main result through an application of the so called *ECO* method formalized by Pinzani et al., see [2].

3.1 The cardinality of $sub(\tilde{\mathcal{A}})_n$

The two cryteria $a1$) and $a2$) given in Section 2, which were used to draw binary increasing trees in a unique and well defined way, are not the only possible. Indeed one can consider the cryterion $a2$) plus the dual version of $a1$). To be more precise, if in a tree a node has only one child, then this child is drawn on the *left* of its direct ancestor and not on the right as it was in $a1$).

We will call those tree which are drawn respecting $a2$) and the new condition described above *left oriented* binary increasing tree. Given such a tree we can still apply the procedure ϕ described in Section 2, without considering step 1), to define a permutation which is not, in general, an André one.

In what follows we want to show that a permutation belongs to $sub_n(\tilde{\mathcal{A}})$ if and only if it is the permutation coming from a left oriented binary increasing tree of size n. As a corollary we will have also that $|sub_n(\tilde{\mathcal{A}})| = |\mathcal{B}_n|$.

Proposition 1 *The following hold:*

i) If π comes from a left oriented binary increasing tree of size n, then $\pi \in sub_n(\tilde{\mathcal{A}})$.

ii) If $\pi \in sub_n(\tilde{\mathcal{A}})$, then π comes from a left oriented binary increasing tree of size n.

Proof. To prove i) take the left oriented tree corresponding to π and, to every node with outdegree 1, add a child with label a number greater than n. Let t be the resulting tree. Then $t \in \tilde{\mathcal{B}}$ and $\pi \lhd \phi(t) \in \tilde{\mathcal{A}}$.

To prove ii) consider that $\pi \lhd \sigma \in \tilde{\mathcal{A}}_m$ with $n \leq m$. Let $t \in \tilde{\mathcal{B}}_m$ be the tree associated to σ. Remove from t all the nodes labelled by a number $k > n$. The obtained tree is left oriented and σ comes from t applying steps 2) and 3) of ϕ. $\square$

3.2 The cardinality of $sub_{n,m}(\tilde{\mathcal{A}})$

In the previous section we have shown that permutations in $sub(\tilde{\mathcal{A}})$ corresponds to those permutations coming from left oriented increasing binary trees when we apply steps 2) and 3) of the mapping ϕ. Now we want to refine the bijection described in the proof of Proposition 1 to consider also the set of left to right minima of such permutations. To do so we need the following definition.

Given a binary increasing tree t consider the following path: we start from the root of t and at each step we move to the child with the least label. In Fig. 2 we have depicted a tree t where such a path, which will be called the *min-path* of t, is highlighted.



Figure 2: The min-path of a binary increasing tree.

Next proposition states the correspondance between left to right minima of a permutation in $sub(\tilde{\mathcal{A}})$ and the min-path in the corresponding tree.

Proposition 2 *If* $\pi \in sub(\tilde{\mathcal{A}})$*, then* $L(\pi)$ *consists of the labels belonging to the min-path of* $\phi^{-1}(\pi)$*.*

Proof. Fistly we observe that when a tree is left oriented then its min-path corresponds to the path going from the root to the leftmost leaf.

Let us consider $t = \phi^{-1}(\pi)$, so t is a left oriented binary increasing tree.

If a node x belongs to the min-path of t, then the entries of π which are on the left of x correspond to the nodes belonging to the left subtree of x. Since t is increasing we have that $x \in L(\pi)$.

If a node x does not belong to the min-path of t, then, going from x to the root of t, we will find a node of outdegree 2 such that x does not belong to its left subtree. Let z be the left child of this node and y the right one. Then $z \prec y \preceq x$ which implies $z \prec x$. Moreover, applying the procedure ϕ, the entry z will be placed on the left of x. From this we have that $x \notin L(\pi)$. $\square$

As an example consider the binary increasing tree depicted in Fig. 2 and draw it according to its left orientation. Then the associated permutation is $\pi = (11\,7\,6\,10\,9\,5\,8\,2\,1\,13\,4\,14\,3\,12)$ and $L(\pi) = \{11, 7, 6, 5, 2, 1\}$.

Thanks to Propositions 1 and 2 we can obtain the enumeration of $sub_{n,m}(\tilde{\mathcal{A}})$ counting the number of binary increasing trees with respect to the size and to the length of the min-path.

3.2.1 A bivariate generating function for $\mathcal{B}$

In what follows we will present a recursive construction for binary increasing trees. In particular we will construct each tree belonging to $\mathcal{B}_{n+1}$ by adding a new node to a tree in $\mathcal{B}_n$. This construction, denoted by Θ, will then be translated into a functional equation. Solving the



Figure 3: First levels of the generating tree associated to Θ.

equation we will obtain a multivariate exponential generating function counting binary increasing trees with respect to size and to the length of the min-path.

Given a tree $t \in \mathcal{B}_n$, Θ simply adds the node labelled '$n+1$' as a child of a node of t having outdegree less than two. More precisely, if $o(t)$ (resp. $p(t)$) is the number of nodes with outdegree 0 (resp. 1) in t, then Θ applied to t produces $o(t)+p(t)$ elements of $\mathcal{B}_{n+1}$ each time adding the new node labelled $n+1$ as a child of the nodes counted in $o(t)+p(t)$. In Fig. 3 the first levels of the generating tree resulting from Θ are depicted.

Note that, if $q(t)$ is the number of nodes with outdegree two in a tree $t \in \mathcal{B}_n$, then $o(t) = q(t)+1$ and $o(t)+p(t)+q(t) = n$. From these relations we have in particular that $p(t) = n - 2o(t) + 1$. The construction Θ can be translated into the following *succession rule* (see [1] [2]) where each tree is represented by the values of its parameters o, l and n, where l represents the length of the min-path (i.e. $l = |L(\phi(t))|$) while n corresponds to the size.

$$(o,l,n) \rightarrow (o,l,n+1)^{o-1}\,(o,l+1,n+1)\,(o+1,l,n+1)^{n-2o+1}.$$

In particular, given a tree t with parameters $o = o(t)$ and $n = n(t)$, the application of Θ on t produces o new trees having size $n+1$ and the same value for o and $n-2o+1$ new trees having both the size and the number of nodes with outdegree zero augmented by one. The starting point of the construction is the unique tree of size one which is represented by the label $(1,1,1)$.

Now consider the exponential generating function

$$F(x,y,z) = \sum_{t \in \mathcal{B}} \frac{x^{o(t)} y^{l(t)} z^{n(t)}}{n(t)\,!},$$

the previous succession rule can be translated as follows into an equation:

$$
\begin{aligned}
F(x,y,z) &= xyz \\
&\quad + \sum_{x^o y^l z^n \in \mathcal{B}} \frac{(o-1)x^o y^l z^{n+1}}{n+1\,!} \\
&\quad + \sum_{x^o y^l z^n \in \mathcal{B}} \frac{x^o y^{l+1} z^{n+1}}{n+1\,!} \\
&\quad + \sum_{x^o y^l z^n \in \mathcal{B}} \frac{(n-2o+1)(x^{o+1} y^l z^{n+1})}{n+1\,!} \\
&= xyz \\
&\quad + x(1-2x) \sum_{x^o y^l z^n \in \mathcal{B}} \frac{o x^{o-1} y^l z^{n+1}}{n+1\,!} \\
&\quad + (y-1) \sum_{x^o y^l z^n \in \mathcal{B}} \frac{x^o y^l z^{n+1}}{n+1\,!} \\
&\quad + xz\, F(x,y,z).
\end{aligned}
$$

From the previous equation we obtain that

$$
\begin{aligned}
F(x,y,z)(1-xz) - xyz &= x(1-2x) \sum_{x^o y^l z^n \in \mathcal{B}} \frac{o x^{o-1} y^l z^{n+1}}{n+1\,!} \\
&\quad + (y-1) \sum_{x^o y^l z^n \in \mathcal{B}} \frac{x^o y^l z^{n+1}}{n+1\,!},
\end{aligned}
$$

differentiating both sides with respect to the variable z we have

$$
F(x,y,z)(1-x-y) - xy = x(1-2x)\frac{\partial F}{\partial x}(x,y,z) + (xz-1)\frac{\partial F}{\partial z}(x,y,z).
$$

The boundary condition to the previous first order partial differential equation is given by

$$
F(x,y,0) = 0.
$$

One can find the desired solution applying the method of *characteristics* or using some of the Mathemathica tools. In the last case, after performing the substitution $x = 1$, one finds that, when $n \geq 2$ and $2 \leq m \leq n$,

$$
|sub_{n,m}(\tilde{\mathcal{A}})| = [y^m] \left(\frac{\partial^n G}{\partial z^n} \right)_{z=0},
$$

where

$$G(y,z) = 2^y \left(\frac{ie^{iz}}{(i+e^{iz})^2} \right)^y \left((-1)^{-\frac{1}{2}+y} \right) H \left[y, \frac{1}{2}+y; 1+y; \frac{4ie^{iz}}{(i+e^{iz})^2} \right]$$

and H is the Gaussian hypergeometric function, see [6] for more details.

The first terms of $|sub_{n,m}(\tilde{\mathcal{A}})|$ are given by the following table.

n/m	2	3	4	5	6	7	8	9	10
2	1	0	0	0	0	0	0	0	0
3	1	1	0	0	0	0	0	0	0
4	1	3	1	0	0	0	0	0	0
5	2	7	6	1	0	0	0	0	0
6	5	20	25	10	1	0	0	0	0
7	16	70	105	65	15	1	0	0	0
8	61	287	490	385	140	21	1	0	0
9	272	1356	2548	2345	1120	266	28	1	0
10	1385	7248	14698	15204	8715	2772	462	36	1

4 Further works

We would like to extend our statistics considering also left to right maxima, right to left maxima and right to left minima of the permutations in $sub(\tilde{\mathcal{A}})$.

References

[1] C. Banderier, M. Bousquet-Melou, A. Denise, P. Flajolet, D. Gardy, and D. Gouyou-Beauchamps. *Generating functions for generating trees.* In: Proceedings of 11-th formal power series and algebraic combinatorics, 40-52 (1999).

[2] E. Barcucci, A. Del Lungo, E. Pergola, R. Pinzani, *ECO: a methodology for the enumeration of combinatorial objects*, Journal of Difference Equations and Applications, Vol. 5 (1999) 435-490.

[3] F. Bergeron, P. Flajolet, B. Salvy, *Varieties of increasing trees.* In J.-C. Raoult, editor, CAAP'92, volume 581 of Lecture Notes in Computer Science, pages 24-48, 1992. Proceedings of the 17th Colloquium on Trees in Algebra and Programming, Rennes, France, February (1992).

[4] C-O. Chow, W. C. Shiu, *Counting Simsun permutations by Descents*, Ann. Comb. 15 (2011) 625-635.

[5] D. Foata, M. P. Schutzenberger, *Nombres d'Euler et permutations alternantes*, In: Bose, R.C. (Ed.) A survey of combinatorial theory, pp. 173-187, North-Holland, Amsterdam (1973).

[6] R. L. Graham, D. Knuth, O. Patashnik, *Concrete Mathematics*, Addison-Wesley (1989).

[7] N. J. A. Sloane, *The On-Line Encyclopedia of Integer Sequences*, available at: http://oeis.org/.